

# A Non-Renormalization Theorem in Three Dimensional Quantum Field Theory

Tomer Shacham\*

Racah Institute of Physics, The Hebrew University of Jerusalem,  
Jerusalem, 91904, Israel

We discuss the two-point functions of the  $U(1)$  current and energy-momentum tensor in certain gapped three-dimensional field theories. We show that the parity-odd part in both of these correlation functions is one-loop exact. In particular, we find a new and simplified derivation of the Coleman-Hill theorem that also clarifies several subtleties in the original argument. For the energy momentum tensor, our result means that the gravitational Chern-Simons term for the background metric is one-loop exact.

## INTRODUCTION

In quantum field theories, the classical description is often modified by quantum corrections; in some cases, these corrections are limited by non-renormalization theorems. In the context of 2+1 dimensional gauge theories [1–4] with massive matter, it was found that the topological mass of the gauge field receives no contributions at two loops [5, 6].<sup>1</sup> Following this discovery, Coleman and Hill proved that no such corrections are possible when the gauge group is Abelian [7].

Their proof deals directly with properties of Feynman graphs. They explicitly show that above one loop, the contribution of each subgroup of graphs with similar topologies is  $\mathcal{O}(\text{momentum}^2)$  and therefore cannot modify the low energy effective action. The reason behind the difference between one and many loops remained implicit in the proof.

Here, we re-derive the Coleman-Hill theorem as a specific case of a more general theorem, and provide a natural explanation for the distinction of the one loop graph. Our argument is as follows. Since in Abelian theories  $A_\mu$  itself is neutral, the parity-odd part (at zero momentum) of  $\langle A_\mu A_\nu \rangle$  is the same as for a background gauge field which can couple to the global  $U(1)$  current,  $j_\mu$ . We argue that if  $\langle j_\mu j_\nu O \rangle$  is  $\mathcal{O}(\text{momentum}^2)$  for an arbitrary scalar  $O$ , so is the rest of the perturbative expansion. We then show that this must be the case for  $\langle j_\mu j_\nu O \rangle$  to satisfy the corresponding Ward identity. Finally, we trace the reason behind this non-renormalization to the lack of gauge invariance of the Chern-Simons Lagrangian.

A closely related statement regarding the non-renormalization of the parity-odd part of  $\langle T_{\mu\nu} T_{\rho\sigma} \rangle$  at zero momentum is proven using similar arguments. The discussion presented here follows the spurion analysis of [8], where it was pointed out that this correlation function does in fact receive higher loop corrections when the gauge group is non-Abelian, due to the topological structure of the theory [9]. Our proof sheds some light on this issue, and it is further investigated in [10] by a perturbative analysis.

## CONSERVED CURRENTS IN THREE DIMENSIONS

A generic QFT can be described by an action  $S = \int \mathcal{L}$  with

$$\mathcal{L} = \mathcal{L}_0 + \sum_i \lambda_i O_i, \quad (1)$$

where  $\lambda_i$  are couplings and  $O_i$  are scalar operators.<sup>2</sup>

We restrict the discussion to cases where  $\mathcal{L}_0$  describes a massive theory which admits to a manifestly Lorentz invariant quantization. In particular, we study three dimensional Abelian gauge theories,  $\text{QED}_3$ , with arbitrary massive matter.<sup>3</sup> For this class of theories,

$$\mathcal{L}_0 = \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{matter}} \quad (2)$$

where

$$\mathcal{L}_{\text{gauge}} = \frac{1}{4} F_{\mu\nu} \left( \frac{1}{e^2} F^{\mu\nu} - i\kappa \epsilon^{\mu\nu\rho} A_\rho \right) - \frac{1}{2\xi e^2} (\partial_\mu A^\mu)^2 \quad (3)$$

describes a gauge field with mass  $\kappa/e^2$ .<sup>4</sup>

We begin with a brief review of Ward identities, in order to establish notation.

### A brief review of Ward identities

When a theory is invariant under a global transformation with a continuous parameter  $\epsilon$ , Noether's theorem guarantees that there exists a classically conserved current, such that

$$\frac{\delta}{\delta\epsilon(x)} \mathcal{S}' + \partial^\mu j_\mu(x) = 0, \quad (4)$$

where the  $'$  denotes a transformed object. If the symmetry is not anomalous (i.e. it survives quantization), correlation functions are independent of the variation. In particular,

$$\frac{\delta}{\delta\epsilon(x)} \langle j_\nu(y) O(z) \rangle' = 0, \quad (5)$$

\* tomer.shacham@phys.huji.ac.il

<sup>1</sup> The topological mass contributes a parity-odd term to the gauge field propagator. At zero momentum, this term is the Chern-Simons coefficient in the effective action.

<sup>2</sup> The subscript 0 will henceforth denote the limit  $\lambda_i \rightarrow 0$ , which corresponds to the free theory.

<sup>3</sup> Even though the non-Abelian theory contains no massless asymptotic states, it does not respect our restriction due to the need to add (massless) ghosts during quantization.

<sup>4</sup> Throughout this paper, we use a Euclidean metric.

and so

$$\frac{\partial}{\partial x_\mu} \langle j_\mu(x) j_\nu(y) O(z) \rangle = - \langle \frac{\delta}{\delta \epsilon(x)} j'_\nu(y) O(z) \rangle - \langle j_\nu(y) \frac{\delta}{\delta \epsilon(x)} O'(z) \rangle. \quad (6)$$

We refer to (6) as the Ward identity for  $\langle j_\mu j_\nu O \rangle$ .

### The $U(1)$ current

Here we take  $j_\mu$  to be the current associated with a global  $U(1)$  symmetry, which acts on the charged fields as

$$\Phi \rightarrow \Phi' = e^{ie\epsilon} \Phi, \quad (7)$$

with  $e$  the “electric” charge.

In a massive theory, the most general possible tensor structure, or parameterization, of  $\langle j_\mu(p) j_\nu(-p) \rangle$  reads

$$a \delta_{\mu\nu} + \delta \kappa \epsilon_{\mu\nu\rho} p^\rho + \mathcal{O}(\text{momentum}^2). \quad (8)$$

Note that only terms in the perturbative expansion of  $\langle j_\mu(p) j_\nu(-p) \rangle$  that are  $\mathcal{O}(\text{momentum})$  can possibly contribute to  $\delta \kappa$ . As advocated, we argue that no such quantum corrections exist. The basic observation is that the entire perturbative expansion around the free theory consists of insertions of the scalar operators  $O_i$  at zero momentum:

$$\begin{aligned} \langle j_\mu(p) j_\nu(-p) \rangle &= \langle j_\mu(p) j_\nu(-p) \rangle_0 - \sum_i \lambda_i \langle j_\mu(p) j_\nu(-p) O_i(0) \rangle_0 \\ &\quad + \sum_{ij} \frac{1}{2} \lambda_i \lambda_j \langle j_\mu(p) j_\nu(-p) O_i(0) O_j(0) \rangle_0 + \dots \end{aligned} \quad (9)$$

In gapped theories, there are no infrared singularities, and so  $\langle j_\mu(p) j_\nu(-p) O_1(0) \dots O_n(0) \rangle$  is well defined as the limit

$$\lim_{k_i \rightarrow 0} \langle j_\mu(p) j_\nu(q) O_1(k_1) \dots O_n(k_n) \rangle, \quad (10)$$

where  $q = -p - \sum_i k_i$ . We can therefore take this limit in two steps,  $k_{i \neq 1} \rightarrow 0$  followed by  $k_1 \rightarrow 0$ .

Consider the parameterization of  $\langle j_\mu(p) j_\nu(q) O_1(k_1) O_2(0) \dots O_n(0) \rangle$ . The insertion of  $O_1(k_1)$  allows  $p$  and  $q$  to be independent; the insertions at zero momentum, on the other hand, do not impose or relax any constraints on the tensor structure.<sup>5</sup> Consequently, this parameterization does not depend on the number of insertions at zero momentum; the different orders in perturbation theory may differ only by the coefficients in the momentum expansion. Therefore, if  $\mathcal{O}(\text{momentum})$  terms in the parameterization of  $\langle j_\mu(p) j_\nu(q) \mathcal{O}(k_1) \rangle$  violate the corresponding Ward identity, they are absent from the rest of the perturbative corrections as well.

As we now show, this is indeed the case. Even though the symmetry group is Abelian, the variation of the current picks up a term

$$\frac{\delta}{\delta \epsilon(x)} \left( i \vec{\phi} \overleftrightarrow{\partial}_\mu \phi \right)'_{(y)} = -2e \vec{\phi} \phi(y) \partial_\mu \delta(x-y), \quad (11)$$

for every charged scalar  $\phi$ , and the Ward identity for  $\langle j_\mu j_\nu O \rangle$  reads

$$\frac{\partial}{\partial x_\mu} \left( \langle j_\mu(x) j_\nu(y) O(z) \rangle - 2e \delta_{\mu\nu} \delta(x-y) \langle \vec{\phi} \phi(y) O(z) \rangle \right) = 0. \quad (12)$$

However, correlation functions are defined up to contact terms; as  $\delta_{\mu\nu} \delta(x-y)$  respects the symmetry of  $j_\mu(x) j_\nu(y)$ , one can absorb the contact term into a redefinition of  $\langle j_\mu j_\nu O \rangle$ . Hence, there exists a regularization scheme in which

$$p^\mu \langle j_\mu(p) j_\nu(q) O(-p-q) \rangle = 0. \quad (13)$$

The parameterization of  $\langle j_\mu(p) j_\nu(q) O(k_1) \rangle$  is given by

$$a' \delta_{\mu\nu} + b \epsilon_{\mu\nu\rho} (p^\rho - q^\rho) + \mathcal{O}(\text{momentum}^2). \quad (14)$$

Since the Ward identity must be satisfied at each order in the momentum expansion, imposing (13) on the parameterization of  $\langle j_\mu(p) j_\nu(-p) \rangle$  gives  $a = 0$ . Similarly, one finds that both  $a'$  and  $b$  vanish;  $\langle j_\mu j_\nu O \rangle$  is then  $\mathcal{O}(\text{momentum}^2)$  for an arbitrary scalar  $O$ . We thus conclude that

$$\delta \kappa = \lim_{p \rightarrow 0} \left( \frac{1}{6} \epsilon^{\mu\nu\lambda} \frac{\partial}{\partial p^\lambda} \langle j_\mu(p) j_\nu(-p) \rangle_0 \right) \quad (15)$$

and does not receive perturbative corrections. As the current is quadratic in the fields, the classical contribution (15) corresponds to a one loop graph.

The reason why the Ward identity prohibits  $\mathcal{O}(\text{momentum})$  terms in  $\langle j_\mu j_\nu O \rangle$  can be understood by coupling the global  $U(1)$  current to a background gauge field  $a_\mu$ , and the deformation  $O_i$  to a background source  $J_i$ . One then defines

$$\langle j_\mu j_\nu O \rangle \equiv \frac{\delta}{\delta a^\mu} \frac{\delta}{\delta a^\nu} \frac{\delta}{\delta J} \mathcal{Z}[a, J_i] \Big|_{a=0, J_i=0} \quad (16)$$

where

$$e^{-\mathcal{Z}[a, J_i]} = \left\langle e^{-\int (j^\mu a_\mu + \sum_i \lambda_i O_i J_i)} \right\rangle. \quad (17)$$

In the language of background fields, the Ward identity corresponds to gauge invariance of the generating functional.

The only term in the derivative expansion of  $\mathcal{Z}[a, J_i]$  with two  $a$ 's and a  $J$  that includes only one derivative is

$$\int d^3x J \epsilon^{\mu\nu\rho} a_\mu \partial_\nu a_\rho. \quad (18)$$

As the Chern-Simons Lagrangian varies by a full derivative under a gauge transformation, (18) is gauge invariant only in the limit of constant  $J$ . In a gapped theory, this limit is well defined, and corresponds to  $\lim_{p \rightarrow 0} \mathcal{O}(p)$ .

<sup>5</sup> Note that relying on the tensor structure is permissible only when the quantization is manifestly Lorentz invariant.

Let us now briefly review the original proof of Coleman and Hill. The basic argument is that subgroups of graphs which contain a propagator of a neutral field cannot contribute to

$$\lim_{p \rightarrow 0} \left( \frac{1}{6} \epsilon^{\mu\nu\lambda} \frac{\partial}{\partial p^\lambda} \langle A_{\mu(p)} A_{\nu(-p)} \rangle \right). \quad (19)$$

A subsequent analysis of the various topologies, excluding the one-loop case, reveals that each subgroup is equivalent to a linear combination of graphs with neutral propagators, and thus do not contribute as well.

We note that since the gauge field itself is neutral, the parity-odd part of its two point function is the same as for a background field; the previous discussion of the one loop exactness of the global  $U(1)$  current captures the same physics and completes the proof. In the language of currents, the one loop contribution is in some sense classical, hence the difference between one and many loops. Moreover, we do not assume minimal coupling; these conclusions hold also for non-trivially improved currents.

### The energy-momentum tensor

Invariance of a theory under the Poincaré group action

$$x^\mu \rightarrow x'^\mu = x^\mu + \epsilon^\mu, \quad (20)$$

implies the conservation of the energy-momentum tensor,

$$\frac{\delta}{\delta \epsilon^\mu(x)} \mathcal{S}' + \partial^\nu T_{\mu\nu}(x) = 0. \quad (21)$$

Note that (21) defines an equivalence class; we shall henceforth refer to  $T_{\mu\nu}$  as the symmetric tensor, which is associated with the coupling to gravity.

In this section, we go through the same procedure of constraining the parameterizations of  $\langle T_{\mu\nu} O \rangle$ ,  $\langle T_{\mu\nu} T_{\rho\sigma} \rangle$ , and  $\langle T_{\mu\nu} T_{\rho\sigma} O \rangle$  by the corresponding Ward identities, which we now derive for completeness.

Under the transformation (20), the fields vary by a Lie derivative with respect to  $\epsilon$ :

$$\Phi' = \Phi + L_\epsilon \Phi. \quad (22)$$

The variation of a scalar field  $\phi$  is

$$\delta\phi = (\epsilon \cdot \partial)\phi, \quad (23)$$

and so the Ward identity for  $\langle T_{\mu\nu} O \rangle$  reads

$$p^\mu \langle T_{\mu\nu(p)} O(q) \rangle = (p+q)_\nu \langle O(p+q) \rangle = 0, \quad (24)$$

where the last equality follows from conservation of momentum. Since  $T_{\mu\nu}$  is symmetric, the parameterization of  $\langle T_{\mu\nu(p)} O(-p) \rangle$  must be proportional to

$$p_\mu p_\nu - p^2 \delta_{\mu\nu} + \mathcal{O}(p^4); \quad (25)$$

taking  $O$  to be the unit operator, one finds that

$$\langle T_{\mu\nu(0)} \rangle = 0. \quad (26)$$

For a symmetric rank 2 tensor,

$$\begin{aligned} \delta T_{\rho\sigma} &= (\epsilon \cdot \partial) T_{\rho\sigma} + (T_{\rho\nu} \partial_\sigma \epsilon^\nu + (\rho \leftrightarrow \sigma)) \\ &= (\epsilon \cdot \partial) T_{\rho\sigma} + \left( \Sigma_{\mu\nu, \rho\sigma}^{(-)\alpha\beta} + \Sigma_{\mu\nu, \rho\sigma}^{(+)\alpha\beta} \right) T_{\alpha\beta} \partial^\mu \epsilon^\nu, \end{aligned} \quad (27)$$

where the second term was decomposed into  $\mu \leftrightarrow \nu$  symmetric and antisymmetric components

$$\Sigma_{\mu\nu, \rho\sigma}^{(\pm)\alpha\beta} \equiv \frac{1}{2} \delta_\rho^\beta (\delta_{\mu\sigma} \delta_\nu^\alpha \pm (\mu \leftrightarrow \nu)) + (\rho \leftrightarrow \sigma). \quad (28)$$

The Ward identity for  $\langle T_{\mu\nu} T_{\rho\sigma} O \rangle$  is given by

$$\begin{aligned} \frac{\partial}{\partial x_\mu} \left( \langle T_{\mu\nu}(x) T_{\rho\sigma}(y) O(z) \rangle + \delta(x-y) \left( \Sigma_{\mu\nu, \rho\sigma}^{(-)\alpha\beta} + \Sigma_{\mu\nu, \rho\sigma}^{(+)\alpha\beta} \right) \langle T_{\alpha\beta}(y) O(z) \rangle \right) \\ = -\delta(x-y) \frac{\partial}{\partial x^\nu} \langle T_{\rho\sigma}(x) O(z) \rangle - \delta(x-y) \frac{\partial}{\partial x^\nu} \langle T_{\rho\sigma}(y) O(x) \rangle. \end{aligned} \quad (29)$$

The contact term containing  $\Sigma^{(\pm)}$  can be absorbed into a redefinition of  $\langle T_{\mu\nu} T_{\rho\sigma} O \rangle$  as it is symmetric in  $\mu \leftrightarrow \nu$ .<sup>6</sup> In momentum space, this reads

$$\begin{aligned} p^\mu \langle T_{\mu\nu(p)} T_{\rho\sigma}(q) O(-p-q) \rangle &= -p^\mu \Sigma_{\mu\nu, \rho\sigma}^{(-)\alpha\beta} \langle T_{\alpha\beta}(p+q) O(-p-q) \rangle \\ &+ q_\sigma \langle T_{\rho\sigma}(q) O(-q) \rangle - (q+p)_\sigma \langle T_{\rho\sigma}(p+q) O(-p-q) \rangle. \end{aligned} \quad (31)$$

Taking  $O$  to be the unit operator and using (26) gives

$$p^\mu \langle T_{\mu\nu(p)} T_{\rho\sigma}(-p) \rangle = 0. \quad (32)$$

The parameterization of  $\langle T_{\mu\nu(p)} T_{\rho\sigma}(-p) \rangle$  is then constrained to

$$\begin{aligned} \dots + \delta\kappa_g \left( (\epsilon_{\mu\rho\lambda} p^\lambda (p_\nu p_\sigma - p^2 \delta_{\nu\sigma}) + (\mu \leftrightarrow \nu)) + \rho \leftrightarrow \sigma \right) \\ + \mathcal{O}(\text{momentum}^4) \end{aligned} \quad (33)$$

where the ellipsis stands for a momentum<sup>2</sup> piece, which plays no role in this discussion; only  $\mathcal{O}(\text{momentum}^3)$  terms in the perturbative expansion can contribute to  $\delta\kappa_g$ . However, such terms in the parameterization of  $\langle T_{\mu\nu(p)} T_{\rho\sigma}(q) O(k_1) \rangle$ ,

$$\begin{aligned} \left( (c_1 (p_\nu q_\sigma + q_\nu p_\sigma) + c_2 (p_\nu p_\sigma + q_\nu q_\sigma) + c_3 \delta_{\nu\sigma} (p^2 + q^2) \right. \\ \left. + c_4 \delta_{\nu\sigma} (p \cdot q)) \epsilon_{\mu\rho\lambda} (p^\lambda - q^\lambda) + (\mu \leftrightarrow \nu) \right) + (\rho \leftrightarrow \sigma), \end{aligned} \quad (34)$$

cannot satisfy the Ward identity (31) as there are no momentum<sup>3</sup> terms in  $\langle T_{\rho\sigma} O \rangle$ . By repeating the arguments of the previous section, we conclude that

$$\delta\kappa_g = \lim_{p \rightarrow 0} \left( \frac{1}{90} \epsilon^{\mu\rho\lambda} \frac{\partial}{\partial p^\lambda} \frac{\partial}{\partial p_\nu} \frac{\partial}{\partial p_\sigma} \langle T_{\mu\nu(p)} T_{\rho\sigma}(-p) \rangle \right). \quad (35)$$

<sup>6</sup> This redefinition is equivalent to taking  $\epsilon_\mu$  to be a Killing vector, i.e.

$$\partial_\mu \epsilon_\nu + \partial_\nu \epsilon_\mu = 0. \quad (30)$$

Although the energy-momentum tensor is not necessarily quadratic in the fields, by considering the Noether current associated with the transformation (22)

$$\epsilon^\nu T_{\mu\nu} = \frac{\partial \mathcal{L}}{\partial (\partial_\mu \Phi)} L_\epsilon \Phi - \epsilon_\mu \mathcal{L}, \quad (36)$$

one finds that only the second term in (36) may not be quadratic, and it clearly cannot contribute to  $\delta\kappa_g$ . The classical contribution (35) therefore corresponds to a one-loop graph.

This result can be equivalently stated in terms of the vacuum polarization tensor of a background graviton, thus making a connection with the  $U(1)$  current discussed in the previous section. The observation that the relevant term in the generating functional is not gauge invariant in the presence of the source  $J$ , translates to a similar statement regarding diffeomorphism non-invariance.

### Acknowledgements

I would like to thank Guy Gur-Ari, Ran Yacoby and especially Shmuel Elitzur and Adam Schwimmer for interesting conversations and useful comments. I am grateful for illuminating discussions with Zohar Komargodski, who also suggested this project. This work was supported in part by the BSF – American-Israel Bi-National Science Foundation,

and by a center of excellence supported by the Israel Science Foundation (grant 1665/10).

- 
- [1] Warren Siegel. Unextended Superfields in Extended Supersymmetry. *Nuclear Physics B*, 156:135, 1979.
  - [2] R Jackiw and S Templeton. How Superrenormalizable Interactions Cure their Infrared Divergences. *Phys.Rev. D*, 23:2291, 1981.
  - [3] S Deser, R Jackiw, and S. Templeton. Topologically Massive Gauge Theories. *Annals of Physics*, 281:409–449, 1982.
  - [4] Jonathan F. Schonfeld. A mass term for three-dimensional gauge fields. *Nuclear Physics B*, 185(1):157–171, July 1981.
  - [5] Yeong-Chuan Kao and Mahiko Suzuki. Radiatively induced topological mass terms in (2+1)-dimensional gauge theories. *Phys. Rev. D*, 31(8):2137–2138, 1985.
  - [6] Marc D. Bernstein and Taejin Lee. Radiative corrections to the topological mass in (2+1)-dimensional electrodynamics. *Phys. Rev. D*, 32(4):1020, 1985.
  - [7] Sidney Coleman and Brian Hill. No more corrections to the Topological Mass Term in QED<sub>3</sub>. *Physics Letters*, 159(B):184–188, 1985.
  - [8] Cyril Closset, Thomas T Dumitrescu, Guido Festuccia, Zohar Komargodski, and Nathan Seiberg. Comments on Chern-Simons Contact Terms in Three Dimensions. *Journal of High Energy Physics*, 91, 2012.
  - [9] Edward Witten. Mathematical Physics Quantum Field Theory and the Jones Polynomial. *Communications in Mathematical Physics*, 121:351–399, 1989.
  - [10] Tomer Shacham. work in progress.